


pmKate2-lyso vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

Location of features

P_{CMV IE}: 1-589
 Enhancer region: 59-465
 TATA box: 554-560
 Transcription start point: 583
 mKate2-LAMP1 fusion
 Start codon (ATG): 626-628
 Lysosomal Associated Membrane Protein 1 (LAMP1) sequence : 626-1846
 Start of mKate2 coding sequence (ATG): 1907-1909
 Stop codon: 2603-2605
 SV40 early mRNA polyadenylation signal
 Polyadenylation signals: 2758-2763 & 2787-2792
 mRNA 3' ends: 2796 & 2808
 f1 single-strand DNA origin: 2855-3310
 Bacterial promoter for expression of Kan^r gene
 -35 region: 3372-3377; -10 region: 3395-3400
 Transcription start point: 3407
 SV40 origin of replication: 3651-3786
 SV40 early promoter
 Enhancer (72-bp tandem repeats): 3484-3555 & 3556-3627
 21-bp repeats: 3631-3651, 3652-3672 & 3674-3694
 Early promoter element: 3707-3713
 Major transcription start points: 3703, 3741, 3747 & 3752
 Kanamycin/neomycin resistance gene
 Neomycin phosphotransferase coding sequences:
 Start codon (ATG): 3835-3837; Stop codon: 4627-4629
 G->A mutation to remove Pst I site: 4017
 C->A (Arg to Ser) mutation to remove BssH II site: 4363
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
 Polyadenylation signals: 4865-4870 & 4878-4883
 pUC plasmid replication origin: 5214-5857

Product	Cat.#	Size
pmKate2-lyso vector	FP312	20 µg
Vector type	mammalian expression vector	
Reporter	mKate2	
Reporter codon usage	mammalian	
Promoter for mKate2	P _{CMV IE}	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	far-red fluorescent labeling of lysosomes	

Vector description

pmKate2-lyso is a mammalian expression vector intended for far-red fluorescent labeling of lysosomes in living cells. The vector encodes far-red fluorescent protein mKate2 targeted to lysosomal membrane by rat Lysosomal Associated Membrane Protein 1 (LAMP-1), fused to the mKate2 N-terminus.

mKate2 codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996].

pmKate2-lyso vector can be used as a source of mKate2-LAMP1 hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

Note: The plasmid DNA was isolated from dam⁺-methylated *E.coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

The vector backbone contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Expression in mammalian cells

pmKate2-lyso vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the mKate2-LAMP1 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

References

- Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143–190.
- Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315–324 / PMID: 8805248

Notice to Purchaser:

mKate2-related materials (also referred to as "Products") are intended for research use only. The Products are covered by U.S. Pat. 7,638,615; European Pat. 1994149; and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>. The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MSDS information is available at <http://www.evrogen.com/MSDS.shtml>