



## Fluorescent sensors for detection of intracellular $\text{Ca}^{2+}$ and $\text{H}_2\text{O}_2$


Evrogen offers genetically-encoded fluorescent sensors for monitoring changes in intracellular concentration of calcium ions (Case12) and hydrogen peroxide (HyPer). Both indicators demonstrate high selectivity and sensitivity of detection, allow precise targeting into various subcellular compartments and real time measuring of signals in natural intracellular surroundings.

Protein	HyPer	Case12
Fluorescence color	green	green
Excitation max, nm	420 and 500	491
Emission max, nm	516	516
Specificity	$\text{H}_2\text{O}_2$	$\text{Ca}^{2+}$
Aggregation	not observed	not observed


### HyPer

Ratiometric sensor for hydrogen peroxide, HyPer, demonstrates submicromolar affinity to  $\text{H}_2\text{O}_2$  and is insensitive to other oxidants tested, like superoxide, oxidized glutathione, nitric oxide, and peroxynitrite. Unlike many chemical sensors, HyPer does not cause artifactual ROS generation and can be used for detection of fast changes of  $\text{H}_2\text{O}_2$  concentration in different cell compartments.


Without  $\text{H}_2\text{O}_2$ , HyPer has two excitation peaks with maxima at 420 nm and 500 nm, and one emission peak with maximum at 516 nm. Upon exposure to  $\text{H}_2\text{O}_2$ , the excitation peak at 420 nm decreases proportionally to the increase in the peak at 500 nm, allowing ratiometric measurement of  $\text{H}_2\text{O}_2$ . Oxidized HyPer can be reduced inside cells.



Ratiometric images of the group of HeLa cells before (A), 20 sec after (B), and 600 sec after (C) addition of 180  $\mu\text{l}$  of  $\text{H}_2\text{O}_2$ . Images were pseudocolored using "ratio" lookup table of NIH ImageJ software: blue-green-red-white colors represent lowest-intermediate-high-highest level of  $\text{H}_2\text{O}_2$ .



HyPer excitation (blue line) and emission (green line) spectra.




Changes in the excitation spectrum of isolated HyPer in response to  $\text{H}_2\text{O}_2$  addition. Emission was measured at 530 nm.


## Case12

High dynamic range sensor for calcium ions, Case12, allows measurement of changes of calcium concentration in a physiological range from hundred nanomoles to micromoles with a high signal-to-noise ratio. Binding of  $\text{Ca}^{2+}$  is fast and reversible, allowing monitoring high-frequency  $\text{Ca}^{2+}$  oscillations. In response to  $\text{Ca}^{2+}$  concentration rise, Case12 shows up to 12-fold increase of fluorescence brightness. Fluorescence of Case12 is characterized by single excitation/emission maxima peaked at 491/516 nm, easily detectable using standard GFP filter sets.

In contrast to conventional calcium sensors, Case12 is stable under physiological pH ( $\text{pK}_a$  7.2 in the presence of  $10 \mu\text{M}$   $\text{Ca}^{2+}$ ). This makes it well suitable for *in vivo* use.




Fluorescence changes of human melanoma-derived M21 cells expressing Case12 in response to  $100 \mu\text{M}$  ATP.




Testing Case12 in living cells. HeLa cells expressing Case12 shown before (A) and after (B) ionophore addition.

For more information, please visit our web-site:  
[www.evrogen.com](http://www.evrogen.com)



Case12 normalized excitation (blue) and emission (green) spectra without  $\text{Ca}^{2+}$  (dotted line) and in the presence of  $1 \text{ mM}$  of  $\text{Ca}^{2+}$  (solid line). Case12 shows multi-fold brightness increase of fluorescence in the response to  $1 \text{ mM}$   $\text{Ca}^{2+}$ .



Dependence of Case12 fluorescence on pH in the presence (solid line) and in the absence (dashed line) of  $\text{Ca}^{2+}$ .

## Available vectors

Vector	Cat#
pCase12-cyto	FP991
pCase12-mem	FP993
pCase12-mito	FP992
pHyPer-cyto	FP941
pHyPer-dMito	FP942
pHyPer-nuc	FP944

For licensing information, please contact Evrogen at [license@evrogen.com](mailto:license@evrogen.com)