

pTagRFP-EB3 vector

Cat# FP365

Vector description

pTagRFP-EB3 is a mammalian expression vector encoding TagRFP-EB3 fusion protein. The vector can be used for fluorescent labeling of EB3 protein in living cells.

TagRFP codon usage is optimized for high expression in mammalian cells, i.e. humanized (Haas *et al.*, 1996). Human microtubule-associated EB3 protein is fused to the TagRFP N-terminus.

pTagRFP-EB3 can be used as a source of TagRFP-EB3 hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.


Note: The plasmid DNA was isolated from dam^+ -methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam^- host and make fresh DNA.

The vector backbone also contains immediate early promoter of cytomegalovirus (P_{CMVIE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3' end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Vector map

For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>



Expression in mammalian cells

pTagRFP-EB3 can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the TagRFP-EB3 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman, 1985].

Location of features

PCMV IE: 1-589
Enhancer region: 59-465
TATA box: 554-560
Transcription start point: 583
EB3: 652-1494
TagRFP: 1516-2229
SV40 early mRNA polyadenylation signal
Polyadenylation signals: 2382-2387 2411-2416
mRNA 3' ends: 2420 2432
f1 single-strand DNA origin: 2479-2934
Bacterial promoter for expression of Kan^r gene
-35 region: 2996-3001
-10 region: 3019-3024
Transcription start point: 3031
SV40 origin of replication: 3275-3410
SV40 early promoter
Enhancer (72-bp tandem repeats): 3108-3179 3180-3251
21-bp repeats: 3255-3275, 3276-3296 3298-3318
Early promoter element: 3331-3337
Major transcription start points: 3327, 3365, 3371 3376
Kanamycin/neomycin resistance gene
Neomycin phosphotransferase coding sequences:
Start codon (ATG): 3459-3461
Stop codon: 4251-4253
G->A mutation to remove Pst I site: 3641
C->A (Arg to Ser) mutation to remove BssH II site: 3987
Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
Polyadenylation signals: 4489-4494 4502-4507
pUC plasmid replication origin: 4838-5481

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

References:

Gorman C. High efficiency gene transfer into mammalian cells. In DNA cloning: A Practical Approach, Vol. II. Ed. D. M. Glover. (IRL Press, Oxford, U.K.). 1985; 143-90.

Haas J, Park EC, Seed B. Codon usage limitation in the expression of HIV-1 envelope glycoprotein. *Curr Biol.* 1996; 6 (3):315-24. / pmid: 8805248

Kozak M. An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res.* 1987; 15 (20):8125-48. / pmid: 3313277

Notice to Purchaser:

Evrogen FP-related products are intended for research use only and covered by Evrogen Patents and/or Patent applications pending. By use of these products, you accept the terms and conditions of the applicable Limited Use Label License (enclosed).

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MATERIAL SAFETY DATA SHEET INFORMATION: To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.