


pHyPer-cyto vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

Location of features

P_{CMV IE}: 1-589
Enhancer region: 59-465
TATA box: 554-560
Transcription start point: 583
HyPer
Kozak consensus translation initiation site: 596-606
Start codon (ATG): 603-605; **Stop codon:** 2037-2039
SV40 early mRNA polyadenylation signal
Polyadenylation signals: 2238-2243 & 2267-2272
mRNA 3' ends: 2276 & 2288
f1 single-strand DNA origin: 2335-2790
Eukaryotic promoter for expression of Kan^r gene
-35 region: 2852-2857; **-10 region:** 2875-2880
Transcription start point: 2887
SV40 origin of replication: 3131-3266
SV40 early promoter
Enhancer (72-bp tandem repeats): 2964-3035 & 3036-3107
21-bp repeats: 3111-3131, 3132-3152 & 3154-3174
Early promoter element: 3187-3193
Major transcription start points: 3183, 3221, 3227 & 3232
Kanamycin/neomycin resistance gene
Neomycin phosphotransferase coding sequences:
Start codon (ATG): 3315-3317; **Stop codon:** 4107-4109
G->A mutation to remove Pst I site: 3497
C->A (Arg to Ser) mutation to remove BssH II site: 3843
Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
Polyadenylation signals: 4345-4350 & 4358-4363
pUC plasmid replication origin: 4694-5337

Product	Cat.#	Size
pHyPer-cyto vector	FP941	20 µg
Vector type	mammalian expression vector	
Reporter	HyPer	
Reporter codon usage	mammalian/E. coli	
Promoter for HyPer	P _{CMV IE}	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	Expression of fluorescent hydrogen peroxide sensor HyPer in mammalian cells under the control of CMV promoter; source of HyPer coding sequence	

Vector description

pHyPer-cyto is a mammalian expression vector encoding a fluorescent sensor HyPer. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of the HyPer sequence [Kozak 1987].

The vector can be also used as a source of HyPer coding sequence. Flanking restriction sites are convenient for excision of HyPer sequence and its further insertion into other expression vectors of choice. Alternatively, HyPer coding sequence can be amplified by PCR.

Note: The plasmid DNA was isolated from dam⁺-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

The vector backbone contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli* and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Expression in mammalian cells

pHyPer-cyto vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive HyPer expression in many cell types. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

References

Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.

Kozak (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / PMID: 3313277

Notice to Purchaser:

The HyPer-related materials (also referred to as "Products") are intended for research use only. Some elements of these materials may be covered by third party patents issued and applicable in certain countries. No license under these patents is conveyed expressly or by implication to the recipient of the materials. Users of these materials may be required to obtain a patent license depending upon the particular application and country in which the materials are received or used.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MATERIAL SAFETY DATA SHEET INFORMATION: To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.