


pPS-CFP2-N vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>

Multiple cloning site (MCS)

```

... G. CTA. GCG. CTA. CCG. GAC. TCA. GAT. CTC. GAG. CTC. AAG. CTT. CGA. ATT. CTG. CAG. TCG. ACG. GTA. CCG. CGG. GCC. CGG. GAT. CCA. CCG. GTC. GCC. ACC. ATG. A . . .
      Nhe I       Bgl II       Sac I       Xho I*       Hind III       Pst I       Kpn I       Apa I       BamH I       PS-CFP2
      * - not unique site.
  
```

Location of features

P_{CMV IE}: 1-589
Enhancer region: 59-465
TATA box: 554-560
Transcription start point: 583
MCS: 591-671
PS-CFP2
Kozak consensus translation initiation site: 672-682
Start codon (ATG): 679-681; **Stop codon:** 1393-1395
SV40 early mRNA polyadenylation signal
Polyadenylation signals: 1549-1554 & 1578-1583
mRNA 3' ends: 1587 & 1599
f1 single-strand DNA origin: 1646-2101
Eukaryotic promoter for expression of Kan^r gene
-35 region: 2163-2168; **-10 region:** 2186-2191
Transcription start point: 2198
SV40 origin of replication: 2442-2577
SV40 early promoter
Enhancer (72-bp tandem repeats): 2275-2346 & 2347-2418
21-bp repeats: 2422-2442, 2443-2463 & 2465-2485
Early promoter element: 2498-2504
Major transcription start points: 2494, 2532, 2538 & 2543
Kanamycin/neomycin resistance gene
Neomycin phosphotransferase coding sequences:
Start codon (ATG): 2626-2628; **Stop codon:** 3418-3420
G->A mutation to remove Pst I site: 2808
C->A (Arg to Ser) mutation to remove BssH II site: 3154
Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
Polyadenylation signals: 3656-3661 & 3669-3674
pUC plasmid replication origin: 4005-4648

References

Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.
 Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248
 Kozak (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

Product	Cat.#	Size
pPS-CFP2-N vector	FP802	20 µg
The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.		
Vector type	mammalian expression vector	
Reporter	PS-CFP2	
Reporter codon usage	mammalian	
Promoter for PS-CFP2	P _{CMV IE}	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	PS-CFP2 expression in mammalian cells; generation of fusions to the PS-CFP2 N-terminus	

Vector description

pPS-CFP2-N is a mammalian expression vector encoding cyan | green fluorescent protein PS-CFP2. The vector allows generation of fusions to the PS-CFP2 N-terminus and expression of PS-CFP2 fusions or PS-CFP2 alone in eukaryotic (mammalian) cells.

PS-CFP2 codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of PS-CFP2 sequence [Kozak 1987]. Multiple cloning site (MCS) is located between P_{CMV IE} and PS-CFP2 coding sequence.

The vector backbone contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli* and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Generation of PS-CFP2-tagged fusions

A localization signal or a gene of interest should be cloned into MCS of the vector. It will be expressed as a fusion to the PS-CFP2 N-terminus when inserted in the same reading frame as PS-CFP2 and no in-frame stop codons are present. The inserted sequence should contain an initiating ATG codon. PS-CFP2-tagged fusions retain fluorescent properties of the native protein allowing fusion localization *in vivo*. Unmodified vector will express PS-CFP2, when transfected into eukaryotic (mammalian) cells.

Note: The plasmid DNA was isolated from dam⁺-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

Expression in mammalian cells

pPS-CFP2-N vector can be transfected into mammalian cells by any known transfection method. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

Notice to Purchaser:

Photoactivatable FP-related products are intended to be used by academic (non-commercial) entities and for research purposes only. Any use of the proprietary nucleic acid or protein other than for research use or by a commercial entity is strictly prohibited. Transfer of this product by purchaser to any other party is specifically prohibited.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MATERIAL SAFETY DATA SHEET INFORMATION: To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.