


## pTagGFP2-mito vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>

### Location of features

P<sub>CMV</sub> IE: 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 TagGFP2-mito fusion  
 Start codon (ATG): 597-599  
 Mitochondrial targeting sequence (MTS): 597-683  
 Start of TagGFP2 coding sequence (ATG): 705-707  
 Stop codon: 1419-1421  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 1574-1579 & 1603-1608  
 mRNA 3' ends: 1612 & 1624  
 f1 single-strand DNA origin: 1671-2126  
 Bacterial promoter for expression of Kan<sup>r</sup> gene  
 -35 region: 2188-2193; -10 region: 2211-2216  
 Transcription start point: 2223  
 SV40 origin of replication: 2467-2602  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 2300-2371 & 2372-2443  
 21-bp repeats: 2447-2467, 2468-2488 & 2490-2510  
 Early promoter element: 2523-2529  
 Major transcription start points: 2519, 2557, 2563 & 2568  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 2651-2653; Stop codon: 3443-3445  
 G->A mutation to remove Pst I site: 2833  
 C->A (Arg to Ser) mutation to remove BssH II site: 3179  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 3681-3686 & 3694-3699  
 pUC plasmid replication origin: 4030-4673

### References

Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-90.

Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-24 / pmid: 8805248

Rizzuto et al. (1989) "A gene specifying subunit VIII of human cytochrome c oxidase is localized to chromosome 11 and is expressed in both muscle and non-muscle tissues." *J Biol Chem*, 264 (18): 10595-600 / pmid: 2543673

Rizzuto et al. (1995) "Chimeric green fluorescent protein as a tool for visualizing subcellular organelles in living cells." *Curr Biol*, 5 (6): 635-42 / pmid: 7552174

Product	Cat.#	Size
pTagGFP2-mito vector	<b>FP197</b>	20 µg

The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.

Vector type	mammalian expression vector
Reporter	TagGFP2
Reporter codon usage	mammalian
Promoter for TagGFP2	P <sub>CMV</sub> IE
Host cells	mammalian
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori
Use	green fluorescent labeling of mitochondria

### Vector description

pTagGFP2-mito is a mammalian expression vector intended for green fluorescent labeling of mitochondria in living cells. The vector encodes green fluorescent protein TagGFP2 fused to mitochondrial targeting sequence (MTS) derived from the subunit VIII of human cytochrome C oxidase [Rizzuto et al. 1989; Rizzuto et al. 1995]. MTS is fused to the TagGFP2 N-terminus.

TagGFP2 codon usage is optimized for high expression in mammalian cells, i.e. humanized [Haas et al. 1996]. pTagGFP2-mito can be used as a source of TagGFP2-MTS hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

The vector backbone contains immediate early promoter of cytomegalovirus (P<sub>CMV</sub> IE) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli* and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Expression in mammalian cells

pTagGFP2-mito can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the TagGFP2-MTS fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

### Notice to Purchaser:

Evrogen Fluorescent Protein Products (the Products) are intended for research use only. The Products are covered by U.S. Pat. 7,417,131 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MATERIAL SAFETY DATA SHEET INFORMATION:** To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.