


pTagRFP-Cx43 vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>

Location of features

PCMV IE: 1-589
 Enhancer region: 59-465
 TATA box: 554-560
 Transcription start point: 583
 Connexin 43: 824-1969
 TagRFP: 1991-2704
 SV40 early mRNA polyadenylation signal
 Polyadenylation signals: 2857-2862 & 2886-2891
 mRNA 3' ends: 2895 & 2907
 f1 single-strand DNA origin: 2954-3409
 Bacterial promoter for expression of Kan^r gene
 -35 region: 3471-3476; -10 region: 3494-3499
 Transcription start point: 3506
 SV40 origin of replication: 3750-3885
 SV40 early promoter
 Enhancer (72-bp tandem repeats): 3583-3654 & 3655-3726
 21-bp repeats: 3730-3750, 3751-3771 & 3773-3793
 Early promoter element: 3806-3812
 Major transcription start points: 3802, 3840, 3846 & 3851
 Kanamycin/neomycin resistance gene
 Neomycin phosphotransferase coding sequences:
 Start codon (ATG): 3934-3936; Stop codon: 4726-4728
 G->A mutation to remove Pst I site: 4116
 C->A (Arg to Ser) mutation to remove BssH II site: 4462
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
 Polyadenylation signals: 4964-4969 & 4977-4982
 pUC plasmid replication origin: 5313-5956

Product	Cat.#	Size
pTagRFP-Cx43 vector	FP364	20 µg

The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.

Vector type	mammalian expression vector
Reporter	TagRFP
Reporter codon usage	mammalian
Promoter for TagRFP	P _{CMV IE}
Host cells	mammalian
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori
Use	red (orange) fluorescent labeling of connexin 43

Vector description

pTagRFP-Cx43 is a mammalian expression vector encoding TagRFP-Cx43 fusion protein. The vector can be used for fluorescent labeling of connexin 43 in living cells.

TagRFP codon usage is optimized for high expression in mammalian cells, i.e. humanized (Haas et al. 1996). Rat connexin 43 is fused to the TagRFP N-terminus.

pTagRFP-Cx43 can be used as a source of TagRFP-Cx43 hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

Note: The plasmid DNA was isolated from dam⁺-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

The vector backbone also contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3' end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Expression in mammalian cells

pTagRFP-Cx43 can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the TagRFP-Cx43 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

References

Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.

Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248

Notice to Purchaser:

Evrogen Fluorescent Protein Products (the Products) are intended for research use only. The Products are covered by U.S. Pat. 7,417,131 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MATERIAL SAFETY DATA SHEET INFORMATION: To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.